

Practice 4.2

Practice sheets are not assessed. The intention is to use material from lectures in preparation for Competence tests and Assignments. You are encouraged to use `thing.[TAB]` and `thing?` in IPython.

Suggested imports:

```
import sympy as sp
x,y,z = sp.symbols('x y z')
#sp.init_printing(use_unicode=True) # uncomment if you like this style
```

When defining a symbol, we can specify things about it:

```
x = sp.symbols('x', real=True) # tells Python that x is a real value i.e. not complex
x = sp.symbols('x', positive=True) # tells Python that x > 0
```

In `sympy` and `numpy`, infinity is represented by two letter "o" s i.e. `sp.oo`

The `sympy` function `sp.N` can be used to convert an expression into a numeric value, e.g.

```
a = sp.sqrt(sp.pi)
a, sp.N(a)
```

```
(sqrt(pi), 1.77245385090552)
```

Classes

Make a Python `class` to represent a cat. Initialise the object with whatever properties you think are important about a cat (perhaps including some sensible defaults), and be sure to include the mass.

Make `__str__` display a human-readable description of the cat: if the mass is greater than 100 kg, tell the user it's a lion; if the mass is less than 10^{-20} kg, tell the user that this is Schrödinger's cat.

Make a list of cats of varying masses.

Symbolic computing

Find the value of the integral $\int_0^1 \cos(-x^2) dx$.

Find the maximum of

$$y = x \exp[-x^2].$$

Hint: Make an expression for y ; differentiate with respect to x ; solve for x for this equal to zero.

Find the normalisation constant for

$$\psi(x) \propto \exp\left[-\frac{m\omega}{2\hbar}x^2\right].$$

Hint: Tell Python that m , \hbar , and ω are positive quantities. Integrate from $-\infty$ to $+\infty$.